

# **Genetic Epidemiology of Breast and Ovarian Cancer**

**Northern California Cancer Center  
and  
Stanford University School of Medicine**

# Prevalence (%) of BRCA1 Mutation Carriers among Population-based nonHispanic White Breast and Ovarian Cancer Patients

	Breast Cancer Patients	Ovarian Cancer Patients
All patients	2.4	9.6
Ashkenazi	5.4	20.6
NonAshkenazi	2.2	8.4
Age (yr)		
<35	7.6	8.3
35-59	6.6	18.9
50-65	0.4	4.4
Family history		
Positive	3.2	19.2
Negative	2.1	7.0

# Prevalence of BRCA1 Mutation Carriers in US NonHispanic White Population (Whittemore et al., CEBP 2004)

	Population Size <sup>a</sup>	No. Carriers <sup>b</sup>	Prevalence <sup>c</sup>
Ashkenazim	3,702,400	44,307	1/84
NonAshkenazim	207,758,226	506,206	1/410
Total	211,460,626	550,513	1/384

a) Estimated from 2000 US Census


b) Estimated using prevalence in cases and penetrance estimates of Antoniou et al. (2003)

c) Assuming 90% test sensitivity.


# Risk in relatives of patients with breast and ovarian cancer

- Jennifer Lee, MD, SUSM
- Esther John, PhD, NCCC
- Valerie McGuire PhD, SUSM
- Dee West, PhD, NCCC, SUSM
- Alice S. Whittemore, PhD, SUSM

### Breast cancer risk in relatives



Ovarian cancer risk in relatives



# Conclusions

- **Additional unidentified genes modify breast & ovarian cancer risks in carriers of BRCA mutations**

# Conclusions

- Unidentified genes affect breast & ovarian cancer risks in noncarriers of BRCA mutations
- The evidence is particularly strong in relatives of early-onset breast cancer patients